

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



Figure 1. Genetic and physical map of recombinant plasmids pHUR849, pDAL201B, pDAL210B, and pDAL200A. The locations of the *pap* genes shown as open bars. Restriction site abbreviations: A, Apa I; B, Bam HI; C, Cla I; E, Eco RI; K, Kpn I; Sa, Sal I; and S, Sma I.

Figure 2. DNA sequences of pHURe49 (A), pDAL201B (B), pDAL210B (C), and pDAL200A (D) *gfpH* structural genes. The nontranscribed DNA strand for each clone shown. Numbering is from the 5' end. The deduced amino acid sequence for the correct frame is shown below each DNA strand. The first amino acid of the mature protein is +1. Stop codon is marked with asterisks. The restriction sites for Cla I (ATGGTA), and Sma I (CCCGGG), are underlined once, respectively.

Title: IMMUNOGENIC PILI  
PRESENTING FOREIGN PEPTIDES,  
THEIR PRODUCTION AND USE  
Inventor(s): Peter O'Hanley et al.  
DOCKET NO.: 050939/0104

pHUR849	1	ATGAGACTGCGATTCTCTGTTCACATTTTCTTTTGGCTGCTGTTTGTTCATGGTGT	60
pDAL200A			
pDAL201B			
pDAL210B			
pHUR849	61	TTTGGCCGGTCCGTTTCTCCGCCCCGCAATGTCCTTCCTGAATACTGGGGAGAAGAGCAC	120
pDAL200A			
pDAL201B			
pDAL210B			
pHUR849	121	GTATGGTGGACGGCAGGGCTGCTTTTCATGGTGAGGTGTCAGACCTGCCTGTACTCTG	180
pDAL200A			
pDAL201B			
pDAL210B			
pHUR849	181	CGCATGGAAGACGCTCGGCAGATTATTGATATGGGGGAAACCCCGGTACGGGATTACAG	240
pDAL200A			
pDAL201B			
pDAL210B			
pHUR849	241	ATTGGTTTCTCGGACCTGAAAGAAAATTCAGCCTCCGGCTCAGGAATTGTGAATTTAAC	300
pDAL200A			
pDAL201B			
pDAL210B			
pHUR849	301	AGTCAGGGTGGGAACCTTTTCTCTGATTCCCGGATAAGGGTGACTTTCGATGGCGTCCGG	360
pDAL200A			
pDAL201B			
pDAL210B			
pHUR849	361	GGTGAAACGCCCGGATAAGTTTAAATTATCCGGTCAGGCAAAAGGCATTAATCTGCAGATA	420
pDAL200A			
pDAL201B			
pDAL210B			
pHUR849	421	GCTGATGTGAGGGGAAATATTGCCCGGGCAGGAAAAGTAATGCCTGCAATACCATTGACG	480
pDAL200A			
pDAL201B			
pDAL210B			
pHUR849	481	GGTAATGAAGAAGCGCTGGATTACCCCTCAGAAATTGTGAGAAACGGAAAAAACTTGAA	540
pDAL200A			
pDAL201B			
pDAL210B			
pHUR849	541	GCCGGAATTATTTTGTCTGTCTGGGATTCGGGTGATTATGAGTGA	588
pDAL200A			
pDAL201B			
pDAL210B			

Figure. 3. Comparison of deduced nucleotide sequence of *papH* genes pHUR849, pDAL200A, pDAL201B and, pDAL210B. The nucleotide(nt) identities, compared with the deduced sequence of *PapH* nt sequence *papH* gene of pHUR849 (upper case), are indicated by blank space, nt differences for the *PapH* genes of pDAL201B, pDAL210B and, pDAL200A, are shown with the corresponding single letter nt code (lower case), respectively. Numbering is from the 5' end. The first nt of coding for the leader sequence is numbered 1, and the first nt coding for the mature protein is numbered 67.

Title: IMMUNOGENIC PILI  
PRESENTING FOREIGN PEPTIDES,  
THEIR PRODUCTION AND USE  
Inventor(s): Peter O'Hanley et al.  
DOCKET NO.: 050939/0104

```

-22                               -1 +1                               18
MRLRFSVPLFFFGCVFVHGVFAcGPFPPPGMSLPEYWGEEH
19
VWWDGRAAFHGEVVRPACTLAMEDAWQIIDMGETPVRDLQ 58
59
NGFSGPERKFSRLRLRNCEFN SQGCNLFSDSRIRVTFDGV R 98
99
GETPDKFNLSCQAKGINLQIADVRGN IARACKVMPAIP L T 138
139
GNEEALDYTLRIVRNKGKKLEAGNYFAVLGFRVDYE 173

```

Figure 4. Comparison of deduced amino acid sequence of *papH* genes pHUR849, pDAL201B, pDAL210B and, pDAL200A. The vertical arrow indicates the postulated cleavage site for the signal peptidase. The amino acid (aa) identities, compared with the deduced sequence of PapH protein of pHUR849 (upper case), are indicated by blank space, aa differences for the PapH proteins of pDAL201B, pDAL210B and pDAL200A, are shown with the corresponding single letter aa code (lower case), respectively. The first aa of the leader sequence is numbered -22, and the first aa of the mature protein is numbered +1.




Figure. 5. Comparison of deduced amino acid sequence of *papH* gene deletion mutants pHUR849-5(*pap-5*), pDAL201B(*pap-21*), pDAL210B(*pap-17*) and, pDAL200A(*pap-200A*). The deduced amino acid sequence of each of the final constructs is shown, (A) pHUR849-5 and, (B) pDAL201B, pDAL210B and, pDAL200A which are identical to each other. The amino acid identities of the proteins are upper case letters. The first amino acid of the mature fusion protein is numbered 1. The underlined sequence indicates the amino acid residues deleted from the mature fusion protein of each strain.